

South West Regional Wound Care Program

My Surgical Wound Self-Care Wound Dressing Guide

Naiiie	Date				
This guide will help you or your caregiver learn how to change your surgical wound dressing					
	to change your dressing: x per week AND whenever the dressing soaks through				
Steps	Instructions				
	Hand sanitizer (70% alcohol) <u>and</u> antibacterial liquid hand soap Gloves				
Step 1	Garbage bag Adhesive remover wipe Tap water or normal saline or sterile water				
	Gauze squares Alcohol wipes Forceps/tweezers Scissors				
Gather your	Cotton tipped applicators/Q-tips Tape Cling				
dressing	Wound Filler (dressing that touches wound base):				
supplies	Wound Covering (the outermost dressing):				
	Other:				
	1. Choose an area to change your dressing that is comfortable for you, has good lighting, and				
	is away from children and pets, and place your dressing supplies there				
Step 2	2. Wash your hands well with warm water and antibacterial liquid soap for at least 15				
Step 2	seconds. Dry your hands using paper towels				
Set up	3. Return to the area you have chosen to change your dressing and open and arrange the				
oct up	garbage bag beside you to put your old dressing and garbage in				
	4. Open the dressing packages that you will need, but take care not to touch the dressings				
	inside. Leave the dressings inside of the packages				
	1. Put on your gloves				
Step 3	2. Carefully remove the old dressing and put it in the garbage bag. If instructed by your nurse,				
	use an adhesive remover wipe to help get the dressing off				
Remove the	3. Remove your gloves and put them in the garbage bag				
old dressing	4. Clean your hands with hand sanitizer. When your hands are dry, put on a new pair of				
	gloves				
	1. Clean the wound with at least 100mL of room temperature:				
	a. Tap water				
	b. Normal saline or sterile water				
	c. Other solution:				
Step 4	Your nurse will instruct you to either:				
	Pour the solution over your wound				
Clean the	Apply the solution using a prefilled bottle				
wound	 Apply the solution using a 30mL syringe and wound irrigating tip 				
	2. Gently pat the skin around the wound dry using gauze				
	3. Apply a barrier wipe to the skin around the wound and allow the skin to air dry				
	NOTE: if you are using a large container of saline or sterile water, date the container when it				
	is first opened and throw out any remaining solution after 24 hours				
Step 5	1. If not already boiled (if you were so instructed by your nurse), clean your forceps/tweezers				
	and scissors well with alcohol swabs and allow them to air dry				
Fill the	2. Using the forceps/tweezers, grasp the wound filler dressing and cut off the amount you				
wound	need to fill the wound. Remove this piece from the dressing package				

	3. Still holding the wound filler dressing with your forceps/tweezers, gently fill the wound with the cut piece of wound filler dressing. Do not fill the wound tight, rather think 'fluffy like a					
	cloud', while making sure to touch all edges of the wound with dressing product. You may					
	need to use a cotton tipped applicator/giant Q tip to help tuck the dressing in place					
Step 6	1. Cover the wound filler with the wound covering dressing as instructed by your nurse					
	2. If instructed by your nurse, tape the dressing in place or apply cling to hold the dressing in					
Cover the	place. If using tape, be careful to not stretch the tape when putting it on or you may cause					
wound	your skin to blister. If using cling, do not wrap it around your foot too tightly or you may cut					
wound	off blood flow to your toes					
	Remove your gloves and put them in the garbage bag. Place any remaining garbage in the					
	garbage bag and close the bag					
	Cleanse your hands with hand sanitizer					
	If you have any leftover wound filler, if instructed by your nurse, place it in a sterile					
	container (like a urine specimen container) using the forceps. Close the container and					
Step 7	write your name, the date and the name of the dressing on the container. In future					
Step /	dressing changes, access the remaining dressing pieces from the container using the					
Clean up	forceps and throw the container and any remaining dressing pieces out after fourteen days					
Clean up	Store all of your dressing supplies in a container with a lid, like a shoe box, banker box, or					
	Rubbermaid container. The container chosen should be put in a clean, dry, safe place,					
	away from the reach of children and pets, like the top of your fridge or a shelf in a closet					
	• If you are running out of supplies, i.e. you have less than a week's supply of dressings left,					
	tell your nurse (if they supply you with the dressings) or visit your pharmacy to purchase					
	more (if you buy your own dressings)					
	Call your visiting nurse (if you have one) or your family doctor or nurse practitioner (if you					
	do not have a visiting nurse) if:					
	 The amount of drainage increases or becomes thick, green/blue or dark yellow/tan 					
Step 8	There is a new or changed wound odor					
	 You develop new or worsening pain in or around the wound 					
When to call	 The wound base develops new or more yellow or black areas/tissue 					
the nurse	 The wound gets bigger or new wounds appear around the original one 					
	 The tissue around the wound appears swollen, red, warm and/or hard to touch 					
	 You develop a fever higher than 38°C (100°F) 					
	 You clean into a tunnel but the same amount of cleansing solution you put in does not 					
	come back out despite several attempts at repositioning					
Additional Notes:						

Nurse's name (or Doctor's name	me if you do not have a visiting nurse):
Nurse's Signature/Status:	
Nurse's Telephone Number:	Date:
· · · · · · · · · · · · · · · · · · ·	